Nightmares: How Trauma-Induced Dreams Worsen Mental Health

AP Seminar

December 2025

Word Count: 1319

Background

For many trauma survivors, nightmares aren't fleeting fears, rather, they are recurring episodes that steadily wear down mental well being. Trauma-induced dreams, often referred to as trauma-related nightmares, are commonly associated with post-traumatic stress disorder (PTSD). With nearly 70% of individuals with psychiatric disorders reporting sleep difficulties, (Mendoza-Alvarez, 2025), it raises the question: What are the negative implications on mental well being from trauma induced dreams? Trauma-induced dreams are consistently associated with impaired mental health, and interfere heavily with recovery for individuals exposed to trauma, contributing to a cycle of distress. According to a clinical studies report conducted by assistant scientists Anthony Reffi and Lia Smith at the International Society for Traumatic Stress Studies, trauma-related nightmares are linked to disrupted Rapid Eye Movement, or REM, sleep patterns and heightened emotional arousal, (Reffi & Smith, 2022). Similarly, the National Library of Medicine reports that persistent trauma-related sleep disturbances are associated with greater PTSD symptom severity, weakening the chance of recovery (Secrist et al. 2019). Together, these findings highlight how trauma-induced dreams perpetuate negative well-being. Inability to gain the necessary benefits of sleep reinforces daytime fatigue, as seen in peer reviewed studies conducted by sleep-health researchers from the National Library of Medicine (Shah et al., 2025), indicating time that is supposed to be dedicated to gaining neurological benefits is being heavily interrupted. Trauma induced nightmares impair mental well-being by disrupting REM sleep and emotional regulation, intensifying PTSD symptoms and self-reinforcing cycles of distress, making comprehensive treatment necessary for effective long-term recovery.

Disrupted Sleep Patterns and Emotional Regulation

The issue of impaired mental health among individuals exposed to trauma exists largely due to the disruption of healthy sleep processes responsible for emotional regulation. Nightmares disrupt REM sleep, impairing emotional regulation. The Sleep Research Society study found that trauma induced dreams interfere with REM sleep, a stage of heightened brain activity involved in emotional processing and memory regulation, reducing the intensity of negative experiences, (Mahr et al. 2025). This increases emotional distress during waking hours. Lack of a functioning system that successfully processes distressing memories causes these disturbances to appear with heightened intensity. This implies sleep disturbances actually help sustain trauma effects. In addition, clinical research conducted from the Journal of Clinical Medicine reports that rather than the nightmares themselves, the distress associated with experiencing nightmares is correlated with severe mental instability. Persistent trauma-related sleep disturbances are associated with emotional reactivity and difficulty regulating emotions, especially among individuals with PTSD, (Akkaoui et al., 2020). Together these findings establish how emotional dysregulation worsens feelings of fear and anxiety surrounding experiencing trauma-induced dreams, further increasing the chances of poor sleep, and therefore mental health.

One promising approach to reducing the mental health impacts of trauma-induced dreams is introducing pharmacological and non-pharmacological interventions designed to alleviate symptoms and reduce nightmare intensity. Clinical research conducted by postdoctoral candidates in the Psychiatry Library Investigation Journal highlights that medications such as prazosin can decrease the frequency and severity of trauma-related nightmares, as it sufficiently antagonizes harmful receptors that stimulate non-REM sleep in the central nervous system, (Paiva et al. 2021). Disrupting these mechanisms in the nervous system can improve sleep

quality, improving daily functioning. While pharmacological approaches target the neurobiological mechanisms that sustain trauma related nightmares, improvements in sleep quality create the conditions necessary for behavioral therapies to address the cycle of emotional disruptance. Complementing this, behavioral health specialists Alyssia Simos and David Berle from the European Journal of Trauma and Dissociation emphasize how cognitive behavioral therapies, including Nightmare Rehearsal Therapy, further reduce nightmares and improve PTSD symptom management (Simos et al., 2023). Building on this, Dr. Ann Wagner and other sleep researchers define Nightmare Rehearsal Therapy as the practice of writing and rehearsing the script of a revised, non-distressing dream (Wagner et al., 2023). Reframing traumatic memories allows the brain to further emphasize less distressing interpretations, reversing stagnant trauma recovery. Ignoring these regulations can trap individuals in a negative cycle, due to the inability to be comfortable with oneself. To combat this, policymakers must balance the accessibility of medication based treatments with the sustained benefits of therapy based care. However, a clinical review from the National Library of Medicine caution that individual trauma histories, comorbid conditions, and treatment adherence can limit outcomes. This indicates that trauma informed, flexible care models are essential to maximize effectiveness, (Malek et al. 2025). Ultimately, it is essential for policymakers to take steps forward towards mitigating the long-term effects of nightmares by implementing medication treatments and therapy based care, in order to provide individuals a path to recovery.

Post-Traumatic Stress Disorder Symptoms

Trauma induced nightmares worsen PTSD symptoms, including intrusive memories, and hyperarousal. A 2025 medical review in PubMed Central reports that persistent trauma induced dreams are associated with increased PTSD symptom severity, as recurrent nightmares sustain

heightened physiological arousal (Sheaves et al. 2023). To corroborate this idea, a 2023 study in the Journal of Clinical Medicine found that individuals who experience frequent trauma related nightmares report higher levels of intrusive thoughts and hyperarousal, or a state of heightened chronic stress, further suggesting sleep disturbances worsen PTSD symptomology through poor sleep quality (Rolling et al., 2023). These studies indicate how trauma-induced dreams are not merely a byproduct of PTSD, but a contributing factor to a greater cycle that intensifies symptom severity and impedes psychological recovery. Without intervention, lack of guided direction could result in increased hopelessness in an attempt to escape from one's own mind.

One proposed solution to the worsening of PTSD symptoms caused by trauma-induced dreams is the targeted treatment of nightmares. Psychiatric researchers emphasize clinical interventions that directly reduce nightmare frequency, arguing that limiting repeated nocturnal exposure to nightmares can alleviate intrusive thoughts and hyperarousal, as hyperarousal "leads to increased chances of subsequent nightmares," (Sheaves et al. 2023). Supporting this perspective, a report from PubMed suggests that trauma induced nightmares function as a reinforcing mechanism for PTSD symptoms, indicating that interventions aimed at disrupting this cycle may improve symptom trajectories, (Reffi et al., 2025). Alleviating PTSD symptoms, present an effective way to shut down this harmful process altogether. In contrast, neuropsychiatric researchers add that focusing solely on nightmare reduction may be insufficient without addressing broader sleep regulation and trauma processing. Peer-reviewed research conducted in Science Direct highlights how integrated treatment approaches combining sleep focused interventions with trauma-informed psychological care are more effective than single method strategies, (Mendoza Alvarez et al. 2025). While nightmare-targeted treatments offer a clear pathway to symptom relief, effectiveness may be limited when underlying sleep

dysregulation, or if unaddressed trauma-induced cognitive patterns remain. This provides an additional perspective by emphasizing the need for more in-depth, comprehensive solutions that address all aspects of the problem. Although resources are required, these findings demonstrate that integrated treatment approaches are most effective. Regardless, it is essential policymakers and officials begin taking steps toward implementing these combined strategies to present a path forward for trauma survivors towards building a new identity.

Conclusion

Trauma-induced nightmares worsen PTSD symptoms and disrupt emotional regulation. Comprehensive treatments must address both sleep and trauma to reduce long term mental health impacts. Targeted interventions can alleviate symptoms, however short term fixes alone do not break the cycle of recurring trauma exposure, highlighting the need for comprehensive recovery strategies. Without intervention trauma survivors will continue to view themselves through the lens of their trauma, instead of who they are beyond the pain.

Works Cited

Akkaoui, M., Lejoyeux, M., Pia d'Ortho, M., & Geoffrey, P. (2020). *Nightmare in Patients with Major Depressive Disorder, Bipolar Disorder, and Psychotic Disorders: A Systematic Review. Journal of Clinical Medicine.* Retrieved from: https://www.mdpi.com/2077-0383/9/12/3990

Mahr, G., Reffi, A., Jankowiak, L., Moore, D., & Drake, C. (2025). *1178 Emotional Dream Content of Acute Trauma Patients: Associations with Interpersonal Violence, Nightmare, and PTSD. Oxford Academic.* Retrieved from: https://academic.oup.com/sleep/article/48/Supplement_1/A509/8134724

Malek, N., Santistevan, A., Yack, L., Kovnick, M., Pracar, S., Berg, E., Metzler, T., Woodward, S., Neylan, T., & Richard, A. (2025). *Evaluating the Relationship Between Emotion Regulation, Mood and Distressing Dreams Using Daily Sleep Diary Reports in Trauma Survivors. National Library of Medicine.* Retrieved from: https://pmc.ncbi.nlm.nih.gov/articles/PMC12426701/

Mendoza-Alvarez, M., Balthasar, Y., Verbraecken, J., Claes, L., van Someren, E., van Marle, H., Vandekerckhove, M., & Picker, L. (2025). *Systemic review: REM sleep, dysphoric dreams and nightmares as transdiagnostic features of psychiatric disorders with emotion dysregulation - Clinical implications. Science Direct.* Retrieved from: https://www.sciencedirect.com/science/article/abs/pii/S1389945724006014

Paiva, H., Filho, I., & Cais, C. (2021). *Using Prazosin to Treat Posttraumatic Stress Disorder and Associations: A Systematic Review Retrieved Psychiatry Investigation Journal.* Retreived from: https://psychiatryinvestigation.org/journal/view.php?doi=10.30773/pi.2020.0411

Reffi, A., Kalmbach, D., Cheng, P., Moore, D., Jennings, M., Mahr, G., Seymour, G., Jankowiak, L., & Drake, C. (2025). *Nightmares and insomnia within the acute aftermath of trauma*

prospectively predict suicidal ideation. National Library of Medicine. Retrieved from: https://pubmed.ncbi.nlm.nih.gov/40265245/

Reffi, A., & Smith, L. (2022). *Sleep, Trauma, and PTSD: Research Brief. International Society for Traumatic Stress Studies.* Retrieved from: https://istss.org/public-resources/friday-fast-facts/sleep-trauma-and-ptsd/

Rolling, J., Rabot, J., Reynaud, E., Kolb, O., Bourgin, P., & Schroder, C. (2023). *Nightmares and Sleep Disturbances in Children with PTSD: A Polysomnographic and Actigraphy Approach Evaluation*. *Journal of Clinical Medicine.* Retrieved from: https://www.mdpi.com/2077-0383/12/20/6570

Secrist, M., John, S., Harper, S., Conners Edge, N., Sigel, B., Sievers, C., & Kramer, T. (2019). *Nightmares in Treatment Seeking Youth: the Role of Cumulative Trauma Exposure. National Library of Medicine.* Retrieved from: https://pmc.ncbi.nlm.nih.gov/articles/PMC7289908/

Simos, A., & Berle, D. (2023). *Phenomenology in Nightmares in PTSD and cPTSD. European Journal of Trauma and Dissociation.* Retrieved from: https://www.sciencedirect.com/science/article/pii/S2468749923000236

Shah, A., Pant, M., Bommasamudram, T., Nayak, K., Roberts, S., Gallagher, C., Vaishali, K., Edwards, B., Tod, D., Davis, F., & Pullinger, S. (2025). *Effects of Sleep Deprivation on Physical and Mental Outcomes: An Umbrella Review. National Library of Medicine.* Retrieved from: https://pmc.ncbi.nlm.nih.gov/articles/PMC12116485/

Sheaves, B., Rek, S., & Freeman, D. (2023). *Nightmares and psychiatric symptoms: A systematic review of longitudinal, experimental, and clinical trial studies. National Library of Medicine.* Retrieved from: https://pmc.ncbi.nlm.nih.gov/articles/PMC10933816/

Wagner, A., Richards, A., Chiros, C., Thuras, P., Parsons, E., Oien, A., Schenck, C., & Irfan, M. *A Retrospective Pilot Study of Imagery Rehearsal Therapy Enhanced with Narrative Therapy Principles for the Treatment of Nightmares in US Military Veterans. National Library of Medicine.* Retrieved from: https://pmc.ncbi.nlm.nih.gov/articles/PMC10773522/